

References

- G. R. Abecasis, S. S. Cherny, W. O. Cookson, and L. R. Cardon. Merlin—rapid analysis of dense genetic maps using sparse gene flow trees. Nat. Genet., 30(1): 97–101, Jan 2002.
- C. Abraham and J. H. Cho. Functional consequences of NOD2 (CARD15) mutations. Inflamm. Bowel Dis., 12(7):641–650, Jul 2006.
- A. Abyzov, A. E. Urban, M. Snyder, and M. Gerstein. CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. Genome Res., 21(6):974–984, Jun 2011.
- I. Akinsheye, A. Alsultan, N. Solovieff, et al. Fetal hemoglobin in sickle cell anemia. Blood, 118(1):19–27, Jul 2011.
- A. Al-Chalabi and C. M. Lewis. Modelling the effects of penetrance and family size on rates of sporadic and familial disease. Hum. Hered., 71(4):281–288, 2011.
- T. J. Albert, M. N. Molla, D. M. Muzny, et al. Direct selection of human genomic loci by microarray hybridization. Nat. Methods, 4(11):903–905, Nov 2007.
- D. Altshuler, J. N. Hirschhorn, M. Klannemark, et al. The common PPARgamma Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. Nat. Genet., 26(1):76–80, Sep 2000.
- D. M. Altshuler, R. A. Gibbs, L. Peltonen, et al. Integrating common and rare genetic variation in diverse human populations. Nature, 467(7311):52–58, Sep 2010.
- C. A. Anderson, G. Boucher, C. W. Lees, et al. Meta-analysis identifies 29 additional ulcerative colitis risk loci, increasing the number of confirmed associations to 47. Nat. Genet., 43(3):246–252, Mar 2011.

- C.A. Anderson, F.H. Pettersson, J.C. Barrett, et al. Evaluating the effects of imputation on the power, coverage, and cost efficiency of genome-wide SNP platforms. The American Journal of Human Genetics, 83(1):112–119, 2008.
- K. Asano, T. Matsushita, J. Umeno, et al. A genome-wide association study identifies three new susceptibility loci for ulcerative colitis in the Japanese population. Nat. Genet., 41(12):1325–1329, Dec 2009.
- O. T. Avery, C. M. Macleod, and M. McCarty. Studies on the Chemical Nature of the Substance Inducing Transformation of Pneumococcal Types: Induction of Transformation by a Desoxyribonucleic Acid Fraction Isolated from Pneumococcus Type II. J. Exp. Med., 79(2):137–158, Feb 1944.
- S.C. Bagley, H. White, and B.A. Golomb. Logistic regression in the medical literature:: Standards for use and reporting, with particular attention to one medical domain. Journal of clinical epidemiology, 54(10):979–985, 2001.
- M. Bahlo, D. R. Booth, S. A. Broadley, et al. Genome-wide association study identifies new multiple sclerosis susceptibility loci on chromosomes 12 and 20. Nat. Genet., 41(7):824–828, Jul 2009.
- S. C. Bain, J. B. Prins, C. M. Hearne, et al. Insulin gene region-encoded susceptibility to type 1 diabetes is not restricted to HLA-DR4-positive individuals. Nat. Genet., 2(3):212–215, Nov 1992.
- J.K. Baker. Trainable grammars for speech recognition. The Journal of the Acoustical Society of America, 65:S132, 1979.
- M. J. Bamshad, S. B. Ng, A. W. Bigham, et al. Exome sequencing as a tool for Mendelian disease gene discovery. Nat. Rev. Genet., 12(11):745–755, Nov 2011.
- L. B. Barreiro, G. Laval, H. Quach, E. Patin, and L. Quintana-Murci. Natural selection has driven population differentiation in modern humans. Nat. Genet., 40(3):340–345, Mar 2008.
- J. C. Barrett and L. R. Cardon. Evaluating coverage of genome-wide association studies. Nat. Genet., 38(6):659–662, Jun 2006.
- J. C. Barrett, D. G. Clayton, P. Concannon, et al. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. Nat. Genet., 41(6):703–707, Jun 2009a.
- J. C. Barrett, S. Hansoul, D. L. Nicolae, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn’s disease. Nat. Genet., 40(8):955–962, Aug 2008.

- J. C. Barrett, J. C. Lee, C. W. Lees, et al. Genome-wide association study of ulcerative colitis identifies three new susceptibility loci, including the HNF4A region. *Nat. Genet.*, 41(12):1330–1334, Dec 2009b.
- T. Barrett, D. B. Troup, S. E. Wilhite, et al. NCBI GEO: archive for functional genomics data sets—10 years on. *Nucleic Acids Res.*, 39(Database issue):D1005–1010, Jan 2011.
- A. Barton, W. Thomson, X. Ke, et al. Rheumatoid arthritis susceptibility loci at chromosomes 10p15, 12q13 and 22q13. *Nature genetics*, 40(10):1156–1159, 2008.
- T. C. Bates, J. F. Price, S. E. Harris, et al. Association of KIBRA and memory. *Neurosci. Lett.*, 458(3):140–143, Jul 2009.
- D. E. Bauer and S. H. Orkin. Update on fetal hemoglobin gene regulation in hemoglobinopathies. *Curr. Opin. Pediatr.*, 23(1):1–8, Feb 2011.
- D. C. Baumgart and S. R. Carding. Inflammatory bowel disease: cause and immunobiology. *Lancet*, 369(9573):1627–1640, May 2007.
- G. W. Beadle and E. L. Tatum. Genetic Control of Biochemical Reactions in *Neurospora*. *Proc. Natl. Acad. Sci. U.S.A.*, 27(11):499–506, Nov 1941.
- A. B. Begovich, V. E. Carlton, L. A. Honigberg, et al. A missense single-nucleotide polymorphism in a gene encoding a protein tyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis. *Am. J. Hum. Genet.*, 75(2):330–337, Aug 2004.
- D. M. Behar, D. Garrigan, M. E. Kaplan, et al. Contrasting patterns of Y chromosome variation in Ashkenazi Jewish and host non-Jewish European populations. *Hum. Genet.*, 114(4):354–365, Mar 2004.
- C.J. Bell, D.L. Dinwiddie, N.A. Miller, et al. Carrier testing for severe childhood recessive diseases by next-generation sequencing. *Sci. Transl. Med.*, 3(65):65ra4, 2011.
- G. I. Bell, S. Horita, and J. H. Karam. A polymorphic locus near the human insulin gene is associated with insulin-dependent diabetes mellitus. *Diabetes*, 33(2):176–183, Feb 1984.
- S. T. Bennett, A. M. Lucassen, S. C. Gough, et al. Susceptibility to human type 1 diabetes at IDDM2 is determined by tandem repeat variation at the insulin gene minisatellite locus. *Nat. Genet.*, 9(3):284–292, Mar 1995.
- J. Berkson. Application of the logistic function to bio-assay. *Journal of the American Statistical Association*, 39(227):357–365, 1944.

- J. Berkson. Why I prefer logits to probits. Biometrics, 7(4):327–339, 1951.
- C.N. Bernstein. The natural history of inflammatory bowel disease. Crohn's Disease and Ulcerative Colitis: From Epidemiology and Immunobiology to a Rational Diagnostic and Therapeutic Approach, page 343, 2011.
- L. G. Biesecker. Opportunities and challenges for the integration of massively parallel genomic sequencing into clinical practice: lessons from the ClinSeq project. Genet. Med., 14(4):393–398, Apr 2012.
- L. G. Biesecker, J. C. Mullikin, F. M. Facio, et al. The ClinSeq Project: piloting large-scale genome sequencing for research in genomic medicine. Genome Res., 19(9):1665–1674, Sep 2009.
- J.M. Bland and D.G. Altman. Statistics notes: the odds ratio. BMJ: British Medical Journal, 320(7247):1468, 2000.
- C.I. Bliss. The calculation of the dosage-mortality curve. Annals of Applied Biology, 22(1):134–167, 1935.
- D. Booth, R. Heard, G. Stewart, et al. Refining genetic associations in multiple sclerosis. Lancet Neurol, 7(7):567–569, Jul 2008.
- F. J. Bosker, C. A. Hartman, I. M. Nolte, et al. Poor replication of candidate genes for major depressive disorder using genome-wide association data. Mol. Psychiatry, 16(5):516–532, May 2011.
- D. Botstein, R. L. White, M. Skolnick, and R. W. Davis. Construction of a genetic linkage map in man using restriction fragment length polymorphisms. Am. J. Hum. Genet., 32(3):314–331, May 1980.
- M. Bradic, J. Costa, and I. M. Chelo. Genotyping with Sequenom. Methods Mol. Biol., 772:193–210, 2011.
- O. J. Brand, J. C. Barrett, M. J. Simmonds, et al. Association of the thyroid stimulating hormone receptor gene (TSHR) with Graves' disease. Hum. Mol. Genet., 18(9):1704–1713, May 2009.
- S. R. Brant. Update on the heritability of inflammatory bowel disease: the importance of twin studies. Inflamm. Bowel Dis., 17(1):1–5, Jan 2011.
- S. M. Bray, J. G. Mulle, A. F. Dodd, et al. Signatures of founder effects, admixture, and selection in the Ashkenazi Jewish population. Proc. Natl. Acad. Sci. U.S.A., 107(37):16222–16227, Sep 2010.
- NE Breslow and BE Storer. General relative risk functions for case-control studies. American journal of epidemiology, 122(1):149–162, 1985.

- M. A. Brown, S. H. Laval, S. Brophy, and A. Calin. Recurrence risk modelling of the genetic susceptibility to ankylosing spondylitis. Ann. Rheum. Dis., 59(11):883–886, Nov 2000.
- B. L. Browning and S. R. Browning. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. Am. J. Hum. Genet., 84(2):210–223, Feb 2009.
- S. R. Browning. Multilocus association mapping using variable-length Markov chains. Am. J. Hum. Genet., 78(6):903–913, Jun 2006.
- S. R. Browning and B. L. Browning. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. Am. J. Hum. Genet., 81(5):1084–1097, Nov 2007.
- S. R. Browning and B. L. Browning. Haplotype phasing: existing methods and new developments. Nat. Rev. Genet., 12(10):703–714, Oct 2011.
- B. Buijsse, R.K. Simmons, S.J. Griffin, and M.B. Schulze. Risk assessment tools for identifying individuals at risk of developing type 2 diabetes. Epidemiol. Rev., 33(1):46–62, 2011.
- P. R. Burton, D. G. Clayton, L. R. Cardon, et al. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. Nat. Genet., 39(11):1329–1337, Nov 2007.
- J. Bustamante, C. Picard, S. Boisson-Dupuis, L. Abel, and J.L. Casanova. Genetic lessons learned from x-linked mendelian susceptibility to mycobacterial diseases. Annals of the New York Academy of Sciences, 1246:92–101, 2011.
- F. Capon, M. J. Bijlmakers, N. Wolf, et al. Identification of ZNF313/RNF114 as a novel psoriasis susceptibility gene. Hum. Mol. Genet., 17(13):1938–1945, Jul 2008.
- A. Cassidy, J.P. Myles, M. van Tongeren, et al. The LLP risk model: an individual risk prediction model for lung cancer. Brit. J. Cancer, 98(2):270–276, 2008.
- N. Chatterjee, J. H. Park, N. Caporaso, and M. H. Gail. Predicting the future of genetic risk prediction. Cancer Epidemiol. Biomarkers Prev., 20(1):3–8, Jan 2011.
- K. Chen, J. W. Wallis, M. D. McLellan, et al. BreakDancer: an algorithm for high-resolution mapping of genomic structural variation. Nat. Methods, 6(9):677–681, Sep 2009.
- S. Chen and G. Parmigiani. Meta-analysis of BRCA1 and BRCA2 penetrance. J. Clin. Oncol., 25(11):1329–1333, Apr 2007.

- W. Chen, D. Stambolian, A. O. Edwards, et al. Genetic variants near TIMP3 and high-density lipoprotein-associated loci influence susceptibility to age-related macular degeneration. Proc. Natl. Acad. Sci. U.S.A., 107:7401–7406, 2010.
- A. G. Clark. Inference of haplotypes from PCR-amplified samples of diploid populations. Mol. Biol. Evol., 7(2):111–122, Mar 1990.
- D. Clayton. Link functions in multi-locus genetic models: implications for testing, prediction, and interpretation. Genet. Epidemiol., 36(4):409–418, May 2012.
- D. Clayton and H. T. Leung. An R package for analysis of whole-genome association studies. Hum. Hered., 64(1):45–51, 2007.
- D. G. Clayton. Prediction and interaction in complex disease genetics: experience in type 1 diabetes. PLoS Genet., 5(7):e1000540, Jul 2009.
- F. S. Collins. Shattuck lecture—medical and societal consequences of the Human Genome Project. N. Engl. J. Med., 341(1):28–37, Jul 1999.
- D. Commenges, H. Jacqmin, L. Letenneur, and C. M. Van Duijn. Score test for familial aggregation in probands studies: application to Alzheimer’s disease. Biometrics, 51(2):542–551, Jun 1995.
- O. Companioni, F. Rodríguez Esparragón, A.M. Fernández-Aceituno, and J.C. Rodríguez Prez. Genetic Variants, Cardiovascular Risk and Genome-Wide Association Studies. Rev. Esp. Cardio., 64(6):509–514, 2011.
- A. Compston and A. Coles. Multiple sclerosis. Lancet, 372(9648):1502–1517, Oct 2008.
- P. Concannon, W. M. Chen, C. Julier, et al. Genome-wide scan for linkage to type 1 diabetes in 2,496 multiplex families from the Type 1 Diabetes Genetics Consortium. Diabetes, 58(4):1018–1022, Apr 2009.
- D. F. Conrad, D. Pinto, R. Redon, et al. Origins and functional impact of copy number variation in the human genome. Nature, 464(7289):704–712, Apr 2010.
- J. D. Cooper, D. J. Smyth, A. M. Smiles, et al. Meta-analysis of genome-wide association study data identifies additional type 1 diabetes risk loci. Nat. Genet., 40(12):1399–1401, Dec 2008.
- J.J. Corneveaux, A.J. Myers, A.N. Allen, et al. Association of CR1, CLU and PICALM with alzheimer’s disease in a cohort of clinically characterized and neuropathologically verified individuals. Hum. Mol. Genet., 19(16):3295–3301, 2010.

- A. Cortes and M. A. Brown. Promise and pitfalls of the Immunochip. Arthritis Res. Ther., 13(1):101, 2011.
- C. Cotsapas, B. F. Voight, E. Rossin, et al. Pervasive sharing of genetic effects in autoimmune disease. PLoS Genet., 7(8):e1002254, Aug 2011.
- J.S. Cramer. The origins and development of the logit model. Logit Models from Economics and Other Fields, pages 149–158, 2003.
- M. P. Creighton, A. W. Cheng, G. G. Welstead, et al. Histone H3K27ac separates active from poised enhancers and predicts developmental state. Proc. Natl. Acad. Sci. U.S.A., 107(50):21931–21936, Dec 2010.
- F. H. Crick, L. Barnett, S. Brenner, and R. J. Watts-Tobin. General nature of the genetic code for proteins. Nature, 192:1227–1232, Dec 1961.
- A. G. Cudworth and H. Festenstein. HLA genetic heterogeneity in diabetes mellitus. Br. Med. Bull., 34(3):285–289, Sep 1978.
- M. J. Daly, J. D. Rioux, S. F. Schaffner, T. J. Hudson, and E. S. Lander. High-resolution haplotype structure in the human genome. Nat. Genet., 29(2):229–232, Oct 2001.
- P. Danoy, K. Pryce, J. Hadler, et al. Association of variants at 1q32 and STAT3 with ankylosing spondylitis suggests genetic overlap with Crohn’s disease. PLoS Genet., 6(12):e1001195, 2010.
- E. V. Davydov, D. L. Goode, M. Sirota, et al. Identifying a high fraction of the human genome to be under selective constraint using GERP++. PLoS Comput. Biol., 6(12):e1001025, 2010.
- P. L. De Jager, X. Jia, J. Wang, et al. Meta-analysis of genome scans and replication identify CD6, IRF8 and TNFRSF1A as new multiple sclerosis susceptibility loci. Nat. Genet., 41(7):776–782, Jul 2009.
- deCODEme. Genetic risk calculation, July 2012. URL <http://demo.decode.me.com/health-watch-information/risk-calculation>.
- A. Deisseroth, A. Nienhuis, J. Lawrence, et al. Chromosomal localization of human beta globin gene on human chromosome 11 in somatic cell hybrids. Proc. Natl. Acad. Sci. U.S.A., 75(3):1456–1460, Mar 1978.
- O. Delaneau, J. Marchini, and J. F. Zagury. A linear complexity phasing method for thousands of genomes. Nat. Methods, 9(2):179–181, Feb 2012.
- A. Dewan, M. Liu, S. Hartman, et al. HTRA1 promoter polymorphism in wet age-related macular degeneration. Science, 314(5801):989–992, Nov 2006.

- H. M. Dick. HLA and disease. Introductory review. Br. Med. Bull., 34(3):271–274, Sep 1978.
- C. Dina, D. Meyre, C. Samson, et al. Comment on "A common genetic variant is associated with adult and childhood obesity". Science, 315(5809):187; author reply 187, Jan 2007.
- A. L. Dixon, L. Liang, M. F. Moffatt, et al. A genome-wide association study of global gene expression. Nat. Genet., 39(10):1202–1207, Oct 2007.
- N. J. Dovichi. DNA sequencing by capillary electrophoresis. Electrophoresis, 18(12-13):2393–2399, Nov 1997.
- I.R. Dowbiggin. Inheriting madness: professionalization and psychiatric knowledge in nineteenth-century France, volume 4. Univ of California Pr on Demand, 1991.
- T. R. Dreszer, D. Karolchik, A. S. Zweig, et al. The UCSC Genome Browser database: extensions and updates 2011. Nucleic Acids Res., 40(Database issue):D918–923, Jan 2012.
- P. C. Dubois, G. Trynka, L. Franke, et al. Multiple common variants for celiac disease influencing immune gene expression. Nat. Genet., 42(4):295–302, Apr 2010.
- R. H. Duerr, K. D. Taylor, S. R. Brant, et al. A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. Science, 314(5804):1461–1463, Dec 2006.
- R. Durbin. Biological sequence analysis: probabilistic models of proteins and nucleic acids. Cambridge Univeristy Press, 1998.
- D. Duricova, N. Pedersen, M. Elkjaer, et al. Overall and cause-specific mortality in Crohn's disease: a meta-analysis of population-based studies. Inflamm. Bowel Dis., 16(2):347–353, Feb 2010.
- R.A. Eeles, Z. Kote-Jarai, A.A. Al Olama, et al. Identification of seven new prostate cancer susceptibility loci through a genome-wide association study. Nat. Genet., 41(10):1116–1121, 2009.
- M. Eisenstein. Oxford nanopore announcement sets sequencing sector abuzz. Nature Biotechnology, 30(4):295–296, 2012.
- R. C. Elston and J. Stewart. A general model for the genetic analysis of pedigree data. Hum. Hered., 21(6):523–542, 1971.
- V. Emilsson, G. Thorleifsson, B. Zhang, et al. Genetics of gene expression and its effect on disease. Nature, 452(7186):423–428, Mar 2008.

- F. Esposito, N. A. Patsopoulos, S. Cepok, et al. IL12A, MPHOSPH9/CDK2AP1 and RGS1 are novel multiple sclerosis susceptibility loci. Genes Immun., 11(5): 397–405, Jul 2010.
- D. M. Evans, C. C. Spencer, J. J. Pointon, et al. Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling in the mechanism for HLA-B27 in disease susceptibility. Nat. Genet., 43(8):761–767, Aug 2011.
- D.S. Falconer. The inheritance of liability to certain diseases, estimated from the incidence among relatives. Annals of Human Genetics, 29(1):51–76, 1965.
- D.S. Falconer and T.F.C Mackay. Introduction to quantitative genetics. Longman., 4th edition, 1996.
- J.B. Fan, A. Oliphant, R. Shen, et al. Highly parallel SNP genotyping. In Cold Spring Harbor symposia on quantitative biology, volume 68, pages 69–78. Cold Spring Harbor Laboratory Press, 2003.
- M. A. Ferreira, M. C. O'Donovan, Y. A. Meng, et al. Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. Nat. Genet., 40(9):1056–1058, Sep 2008.
- R. C. Ferreira, Q. Pan-Hammarstrom, R. R. Graham, et al. Association of IFIH1 and other autoimmunity risk alleles with selective IgA deficiency. Nat. Genet., 42(9):777–780, Sep 2010.
- E. A. Festen, P. Goyette, R. Scott, et al. Genetic variants in the region harbouring IL2/IL21 associated with ulcerative colitis. Gut, 58(6):799–804, Jun 2009.
- O. Filipe-Santos, J. Bustamante, M.H. Haverkamp, et al. X-linked susceptibility to mycobacteria is caused by mutations in NEMO impairing CD40-dependent IL-12 production. The Journal of experimental medicine, 203(7):1745–59, 2006.
- R.A. Fisher. The Correlation Between Relatives on the Supposition of Mendelian Inheritance. Transactions of the Royal Society of Edinburgh, 52:399–433, 1918.
- R.A. Fisher. The detection of linkage with “dominant” abnormalities. Annals of Human Genetics, 6(2):187–201, 1935.
- T. Fiskerstrand, N. Arshad, B. I. Haukanes, et al. Familial diarrhea syndrome caused by an activating GUCY2C mutation. N. Engl. J. Med., 366(17):1586–1595, Apr 2012.
- N. Fleshner. Re: Familial clustering of breast and prostate cancers and risk of postmenopausal breast cancer. Journal of the National Cancer Institute, 87(7): 536–537, 1995.

- J. L. Flynn, J. Chan, K. J. Triebold, et al. An essential role for interferon gamma in resistance to *Mycobacterium tuberculosis* infection. J. Exp. Med., 178(6): 2249–2254, Dec 1993.
- S. P. Fodor, J. L. Read, M. C. Pirrung, et al. Light-directed, spatially addressable parallel chemical synthesis. Science, 251(4995):767–773, Feb 1991.
- A. Fontalba, O. Gutierrez, and J. L. Fernandez-Luna. NLRP2, an inhibitor of the NF-kappaB pathway, is transcriptionally activated by NF-kappaB and exhibits a nonfunctional allelic variant. J. Immunol., 179(12):8519–8524, Dec 2007.
- A. Franke, T. Balschun, T. H. Karlsen, et al. Sequence variants in IL10, ARPC2 and multiple other loci contribute to ulcerative colitis susceptibility. Nat. Genet., 40(11):1319–1323, Nov 2008.
- A. Franke, D. P. McGovern, J. C. Barrett, et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn’s disease susceptibility loci. Nat. Genet., 42(12):1118–1125, Dec 2010.
- K.A. Frazer, D.G. Ballinger, D.R. Cox, et al. A second generation human haplotype map of over 3.1 million SNPs. Nature, 449(7164):851–861, 2007.
- P. Frederic and F. Lad. A technical note on the logitnormal distribution. Research Report, Mathematics and Statistics department at Canterbury University (NZ), 2003.
- J. D. Freeman, R. L. Warren, J. R. Webb, B. H. Nelson, and R. A. Holt. Profiling the T-cell receptor beta-chain repertoire by massively parallel sequencing. Genome Res., 19(10):1817–1824, Oct 2009.
- J. Freudenberg, H. S. Lee, B. G. Han, et al. Genome-wide association study of rheumatoid arthritis in Koreans: population-specific loci as well as overlap with European susceptibility loci. Arthritis Rheum., 63(4):884–893, Apr 2011.
- G. Fu, J. Hu, N. Niederberger-Magnenat, et al. Protein kinase CH is required for T cell activation and homeostatic proliferation. Sci Signal, 4(202):ra84, 2011.
- A.E. Garrod. The incidence of alkaptonuria: a study in chemical individuality. Lancet, 2(24):1616–1620, 1902.
- V. Gateva, J. K. Sandling, G. Hom, et al. A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus. Nat. Genet., 41(11):1228–1233, Nov 2009.
- E. O. Glocker, D. Kotlarz, K. Boztug, et al. Inflammatory bowel disease and mutations affecting the interleukin-10 receptor. N. Engl. J. Med., 361(21):2033–2045, Nov 2009a.

- E.O. Glocker, A. Hennigs, M. Nabavi, et al. A homozygous card9 mutation in a family with susceptibility to fungal infections. The New England journal of medicine, 361(18):1727–35, 2009b.
- A. Goate, M. C. Chartier-Harlin, M. Mullan, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer’s disease. Nature, 349(6311):704–706, Feb 1991.
- A. Gonzalez-Perez and N. Lopez-Bigas. Improving the assessment of the outcome of nonsynonymous SNVs with a consensus deleteriousness score, *Condel*. Am. J. Hum. Genet., 88(4):440–449, Apr 2011.
- E.L. Goode, G. Chenevix-Trench, H. Song, et al. A genome-wide association study identifies susceptibility loci for ovarian cancer at 2q31 and 8q24. Nat. Genet., 42(10):874–879, 2010.
- Y. Goto, A. Hishida, K. Matsuo, et al. PRKCH gene polymorphism is associated with the risk of severe gastric atrophy. Gastric Cancer, 13(2):90–94, Jun 2010.
- D. M. Greenawalt, R. Dobrin, E. Chudin, et al. A survey of the genetics of stomach, liver, and adipose gene expression from a morbidly obese cohort. Genome Res., 21(7):1008–1016, Jul 2011.
- R. C. Griffiths and P. Marjoram. Ancestral inference from samples of DNA sequences with recombination. J. Comput. Biol., 3(4):479–502, 1996.
- W. Guan, A. Pluzhnikov, N. J. Cox, et al. Meta-analysis of 23 type 2 diabetes linkage studies from the International Type 2 Diabetes Linkage Analysis Consortium. Hum. Hered., 66(1):35–49, 2008.
- S. G. Guerra, T. J. Vyse, and D. S. Cunninghame Graham. The genetics of lupus: a functional perspective. Arthritis Res Ther, 14(3):211, May 2012.
- J. Gulcher and K. Stefansson. Genetic risk information for common diseases may indeed be already useful for prevention and early detection. Eur. J. Clin. Invest., 40(1):56–63, Jan 2010.
- DG Haegert. Analysis of the threshold liability model provides new understanding of causation in autoimmune diseases. Medical hypotheses, 63(2):257–261, 2004.
- J.B.S. Haldane. Methods for the detection of autosomal linkage in man. Annals of Human Genetics, 6(1):26–65, 1934.
- W. D. Hall, R. Mathews, and K. I. Morley. Being more realistic about the public health impact of genomic medicine. PLoS Med., 7(10), Oct 2010.

- J. Hampe, S. Schreiber, S. H. Shaw, et al. A genomewide analysis provides evidence for novel linkages in inflammatory bowel disease in a large European cohort. Am. J. Hum. Genet., 64(3):808–816, Mar 1999.
- J. W. Han, H. F. Zheng, Y. Cui, et al. Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus. Nat. Genet., 41(11):1234–1237, Nov 2009.
- R. E. Handsaker, J. M. Korn, J. Nemesh, and S. A. McCarroll. Discovery and genotyping of genome structural polymorphism by sequencing on a population scale. Nat. Genet., 43(3):269–276, Mar 2011.
- J. B. Harley, M. E. Alarcon-Riquelme, L. A. Criswell, et al. Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXX, KIAA1542 and other loci. Nat. Genet., 40(2):204–210, Feb 2008.
- D. Harold, R. Abraham, P. Hollingworth, et al. Genome-wide association study identifies variants at CLU and PICALM associated with alzheimer’s disease. Nat. Genet., 41(10):1088–1093, 2009.
- J. Harrow, F. Denoeud, A. Frankish, et al. GENCODE: producing a reference annotation for ENCODE. Genome Biol., 7 Suppl 1:1–9, 2006.
- M. Heinig, E. Petretto, C. Wallace, et al. A trans-acting locus regulates an antiviral expression network and type 1 diabetes risk. Nature, 467(7314):460–464, Sep 2010.
- A. Herbert, N. P. Gerry, M. B. McQueen, et al. A common genetic variant is associated with adult and childhood obesity. Science, 312(5771):279–283, Apr 2006.
- L. A. Hindorff, P. Sethupathy, H. A. Junkins, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. Proc. Natl. Acad. Sci. U.S.A., 106(23):9362–9367, Jun 2009.
- D. A. Hinds, L. L. Stuve, G. B. Nilsen, et al. Whole-genome patterns of common DNA variation in three human populations. Science, 307(5712):1072–1079, Feb 2005.
- A. D. Hingorani, J. P. Casas, D. I. Swerdlow, et al. The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. Lancet, 379(9822):1214–1224, Mar 2012.
- G. M. Hirschfield, X. Liu, C. Xu, et al. Primary biliary cirrhosis associated with HLA, IL12A, and IL12RB2 variants. N. Engl. J. Med., 360(24):2544–2555, Jun 2009.

- S. M. Holland, F. R. DeLeo, H. Z. Elloumi, et al. STAT3 mutations in the hyper-IgE syndrome. N. Engl. J. Med., 357(16):1608–1619, Oct 2007.
- R.S. Houlston, E. Webb, P. Broderick, et al. Meta-analysis of genome-wide association data identifies four new susceptibility loci for colorectal cancer. Nat. Genet., 40(12):1426–1435, 2008.
- B. Howie, C. Fuchsberger, M. Stephens, J. Marchini, and G. R. Abecasis. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. Nat. Genet., 44(8):955–959, Aug 2012.
- B. Howie, J. Marchini, and M. Stephens. Genotype imputation with thousands of genomes. G3 (Bethesda), 1(6):457–470, Nov 2011.
- B. N. Howie, P. Donnelly, and J. Marchini. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. PLoS Genet., 5(6):e1000529, Jun 2009.
- X. Hu, H. Kim, E. Stahl, et al. Integrating autoimmune risk loci with gene-expression data identifies specific pathogenic immune cell subsets. Am. J. Hum. Genet., 89(4):496–506, Oct 2011.
- J. Huang, D. Ellinghaus, A. Franke, B. Howie, and Y. Li. 1000 Genomes-based imputation identifies novel and refined associations for the Wellcome Trust Case Control Consortium phase 1 Data. Eur. J. Hum. Genet., 20(7):801–805, Jul 2012.
- L. Huang, Y. Li, A.B. Singleton, et al. Genotype-imputation accuracy across worldwide human populations. The American Journal of Human Genetics, 84(2):235–250, 2009.
- J. P. Hugot, M. Chamaillard, H. Zouali, et al. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn’s disease. Nature, 411(6837):599–603, May 2001.
- K. A. Hunt, D. J. Smyth, T. Balschun, et al. Rare and functional SIAE variants are not associated with autoimmune disease risk in up to 66,924 individuals of European ancestry. Nat. Genet., 44(1):3–5, Jan 2012.
- K. A. Hunt, A. Zhernakova, G. Turner, et al. Newly identified genetic risk variants for celiac disease related to the immune response. Nat. Genet., 40(4):395–402, Apr 2008.
- FB Hutt, WR LAMOREUX, et al. Genetics of the fowl. 11. a linkage map for six chromosomes. Journal of Heredity, 31:231–235, 1940.

- G. Hyatt, R. Melamed, R. Park, et al. Gene expression microarrays: glimpses of the immunological genome. *Nat. Immunol.*, 7(7):686–691, Jul 2006.
- M. Imielinski, R. N. Baldassano, A. Griffiths, et al. Common variants at five new loci associated with early-onset inflammatory bowel disease. *Nat. Genet.*, 41(12):1335–1340, Dec 2009.
- V. M. Ingram. Gene mutations in human haemoglobin: the chemical difference between normal and sickle cell haemoglobin. *Nature*, 180(4581):326–328, Aug 1957.
- International HapMap Consortium. The International HapMap Project. *Nature*, 426(6968):789–796, Dec 2003.
- International HapMap Consortium. A haplotype map of the human genome. *Nature*, 437(7063):1299–1320, Oct 2005.
- International Parkinson’s Disease Genomics Consortium and Wellcome Trust Case Control Consortium 2. A two-stage meta-analysis identifies several new Loci for Parkinson’s disease. *PLoS Genet.*, 7:e1002142, 2011.
- J. P. Ioannidis, R. Tarone, and J. K. McLaughlin. The false-positive to false-negative ratio in epidemiologic studies. *Epidemiology*, 22(4):450–456, Jul 2011.
- D. A. Jackson, R. H. Symons, and P. Berg. Biochemical method for inserting new genetic information into DNA of Simian Virus 40: circular SV40 DNA molecules containing lambda phage genes and the galactose operon of *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.*, 69(10):2904–2909, Oct 1972.
- E. Jakkula, V. Leppa, A. M. Sulonen, et al. Genome-wide association study in a high-risk isolate for multiple sclerosis reveals associated variants in *STAT3* gene. *Am. J. Hum. Genet.*, 86(2):285–291, Feb 2010.
- D. Jawaheer, M. F. Seldin, C. I. Amos, et al. Screening the genome for rheumatoid arthritis susceptibility genes: a replication study and combined analysis of 512 multicase families. *Arthritis Rheum.*, 48(4):906–916, Apr 2003.
- G. A. Jervis. Phenylpyruvic oligophrenia deficiency of phenylalanine-oxidizing system. *Proc. Soc. Exp. Biol. Med.*, 82(3):514–515, Mar 1953.
- T. Jess, M. Gamborg, P. Munkholm, and T. I. Sørensen. Overall and cause-specific mortality in ulcerative colitis: meta-analysis of population-based inception cohort studies. *Am. J. Gastroenterol.*, 102(3):609–617, Mar 2007.
- Norman L. Johnson, Samuel Kotz, and N. Balakrishnan. *Continuous Univariate Distributions*, volume I. John Wiley & Sons Inc., New York, 2nd edition, 1994. ISBN 0-471-58495-9.

- L. Jostins, K. I. Morley, and J. C. Barrett. Imputation of low-frequency variants using the HapMap3 benefits from large, diverse reference sets. Eur. J. Hum. Genet., 19(6):662–666, Jun 2011.
- T. M. Karafet, F. L. Mendez, M. B. Meilerman, et al. New binary polymorphisms reshape and increase resolution of the human Y chromosomal haplogroup tree. Genome Res., 18(5):830–838, May 2008.
- A. Kaser, A. H. Lee, A. Franke, et al. XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. Cell, 134(5):743–756, Sep 2008.
- F. K. Kavvoura, T. Akamizu, T. Awata, et al. Cytotoxic T-lymphocyte associated antigen 4 gene polymorphisms and autoimmune thyroid disease: a meta-analysis. J. Clin. Endocrinol. Metab., 92(8):3162–3170, Aug 2007.
- J. Keane, S. Gershon, R. P. Wise, et al. Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. N. Engl. J. Med., 345(15):1098–1104, Oct 2001.
- B. Kerem, J. M. Rommens, J. A. Buchanan, et al. Identification of the cystic fibrosis gene: genetic analysis. Science, 245(4922):1073–1080, Sep 1989.
- JFC Kingman. On the genealogy of large populations. Journal of Applied Probability, pages 27–43, 1982.
- JB Kirsner. Genetic aspects of inflammatory bowel disease. Clin Gastroenterol, 2(557):76, 1973.
- J.B. Kirsner. The historical basis of the idiopathic inflammatory bowel diseases. Inflammatory Bowel Diseases, 1(1):2–26, 1995.
- G. D. Kitsios, N. Tangri, P. J. Castaldi, and J. P. Ioannidis. Laboratory mouse models for the human genome-wide associations. PLoS ONE, 5(11):e13782, 2010.
- R. J. Klein, C. Zeiss, E. Y. Chew, et al. Complement factor H polymorphism in age-related macular degeneration. Science, 308(5720):385–389, Apr 2005.
- R. G. Knowlton, O. Cohen-Haguenauer, N. Van Cong, et al. A polymorphic DNA marker linked to cystic fibrosis is located on chromosome 7. Nature, 318(6044):380–382, 1985.
- A. Kong and N. J. Cox. Allele-sharing models: LOD scores and accurate linkage tests. Am. J. Hum. Genet., 61(5):1179–1188, Nov 1997.

- S. V. Kozyrev, A. K. Abelson, J. Wojcik, et al. Functional variants in the B-cell gene BANK1 are associated with systemic lupus erythematosus. Nat. Genet., 40(2):211–216, Feb 2008.
- P. Kraft and D. J. Hunter. Genetic risk prediction—are we there yet? N. Engl. J. Med., 360(17):1701–1703, Apr 2009.
- L. Kruglyak, M. J. Daly, M. P. Reeve-Daly, and E. S. Lander. Parametric and nonparametric linkage analysis: a unified multipoint approach. Am. J. Hum. Genet., 58(6):1347–1363, Jun 1996.
- S. Kugathasan, R. N. Baldassano, J. P. Bradfield, et al. Loci on 20q13 and 21q22 are associated with pediatric-onset inflammatory bowel disease. Nat. Genet., 40(10):1211–1215, Oct 2008.
- E. Lander and L. Kruglyak. Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. Nat. Genet., 11(3):241–247, Nov 1995.
- E. S. Lander and P. Green. Construction of multilocus genetic linkage maps in humans. Proc. Natl. Acad. Sci. U.S.A., 84(8):2363–2367, Apr 1987.
- E. S. Lander, L. M. Linton, B. Birren, et al. Initial sequencing and analysis of the human genome. Nature, 409(6822):860–921, Feb 2001.
- J.E. Lane-Claypon et al. A further report on cancer of the breast with special reference to its associated antecedent conditions. Ministry of Health. Reports on Public Health and Medical Subjects., 32, 1926.
- T. Langae and M. Ronaghi. Genetic variation analyses by Pyrosequencing. Mutat. Res., 573(1-2):96–102, Jun 2005.
- R. M. Lawn, E. F. Fritsch, R. C. Parker, G. Blake, and T. Maniatis. The isolation and characterization of linked delta- and beta-globin genes from a cloned library of human DNA. Cell, 15(4):1157–1174, Dec 1978.
- S. Q. Le and R. Durbin. SNP detection and genotyping from low-coverage sequencing data on multiple diploid samples. Genome Res., 21(6):952–960, Jun 2011.
- J. Lederberg. Infectious disease as an evolutionary paradigm. Emerging Infect. Dis., 3(4):417–423, 1997.
- J. C. Lee, P. A. Lyons, E. F. McKinney, et al. Gene expression profiling of CD8+ T cells predicts prognosis in patients with Crohn disease and ulcerative colitis. J. Clin. Invest., 121(10):4170–4179, Oct 2011.

- L. G. Lee, C. R. Connell, and W. Bloch. Allelic discrimination by nick-translation PCR with fluorogenic probes. Nucleic Acids Res., 21(16):3761–3766, Aug 1993.
- C. W. Lees, J. C. Barrett, M. Parkes, and J. Satsangi. New IBD genetics: common pathways with other diseases. Gut, 60(12):1739–1753, Dec 2011.
- H. Li and R. Durbin. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics, 25(14):1754–1760, Jul 2009.
- H. Li, B. Handsaker, A. Wysoker, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics, 25(16):2078–2079, Aug 2009.
- J. Li and W.B. Wang. Tag snp selection. Analysis of Complex Disease Association Studies: A Practical Guide, page 49, 2010.
- J. Z. Li, D. M. Absher, H. Tang, et al. Worldwide human relationships inferred from genome-wide patterns of variation. Science, 319(5866):1100–1104, Feb 2008.
- N. Li and M. Stephens. Modeling linkage disequilibrium and identifying recombination hotspots using single-nucleotide polymorphism data. Genetics, 165(4):2213–2233, Dec 2003.
- Y. Li, C. Sidore, H. M. Kang, M. Boehnke, and G. R. Abecasis. Low-coverage sequencing: implications for design of complex trait association studies. Genome Res., 21(6):940–951, Jun 2011.
- Y. Li, C. J. Willer, J. Ding, P. Scheet, and G. R. Abecasis. MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. Genet. Epidemiol., 34(8):816–834, Dec 2010.
- D.C. Link, L.G. Schuettelpelz, D. Shen, et al. Identification of a novel TP53 cancer susceptibility mutation through whole-genome sequencing of a patient with therapy-related AML. JAMA : the journal of the American Medical Association, 305(15):1568–76, 2011.
- P. Linsel-Nitschke, A. Gotz, J. Erdmann, et al. Lifelong reduction of LDL-cholesterol related to a common variant in the LDL-receptor gene decreases the risk of coronary artery disease—a Mendelian Randomisation study. PLoS ONE, 3(8):e2986, 2008.
- R. J. Lipshutz, D. Morris, M. Chee, et al. Using oligonucleotide probe arrays to access genetic diversity. BioTechniques, 19(3):442–447, Sep 1995.
- Q. R. Liu, T. Drgon, C. Johnson, et al. Addiction molecular genetics: 639,401 SNP whole genome association identifies many "cell adhesion" genes. Am. J. Med. Genet. B Neuropsychiatr. Genet., 141B(8):918–925, Dec 2006.

- A. Lleo, L. Moroni, L. Caliari, and P. Invernizzi. Autoimmunity and Turner's syndrome. Autoimmun Rev, 11(6-7):A538–543, May 2012.
- D.M. Lloyd-Jones, K. Liu, L. Tian, and P. Greenland. Narrative review: Assessment of C-reactive protein in risk prediction for cardiovascular disease. Ann. Intern. Med., 145(1):35–42, 2006.
- R. J. Loos, I. Barroso, S. O'rahilly, and N. J. Wareham. Comment on "A common genetic variant is associated with adult and childhood obesity". Science, 315(5809):187; author reply 187, Jan 2007.
- Q. Lu and R. C. Elston. Using the optimal receiver operating characteristic curve to design a predictive genetic test, exemplified with type 2 diabetes. Am. J. Hum. Genet., 82(3):641–651, Mar 2008.
- Luke Jostins. YFitter: a program for assigning haplogroups using maximum likelihood, 2011. URL <http://sourceforge.net/projects/yfitter/>. v0.2.
- V. Lyssenko, A. Jonsson, P. Almgren, et al. Clinical risk factors, DNA variants, and the development of type 2 diabetes. N. Engl. J. Med., 359(21):2220–2232, 2008.
- D. G. MacArthur, S. Balasubramanian, A. Frankish, et al. A systematic survey of loss-of-function variants in human protein-coding genes. Science, 335(6070):823–828, Feb 2012.
- R. Mackelprang, C. S. Carlson, L. Subrahmanyam, et al. Sequence variation in the human T-cell receptor loci. Immunol. Rev., 190:26–39, Dec 2002.
- M. Macpherson, B. Naughton, A. Hsu, and J. Mountain. Estimating genotype-specific incidence for one or several loci, November 2007. URL <https://www.23andme.com/for/scientists/>.
- P. Maerten, C. Shen, S. Colpaert, et al. Involvement of interleukin 18 in Crohn's disease: evidence from in vitro analysis of human gut inflammatory cells and from experimental colitis models. Clin. Exp. Immunol., 135(2):310–317, Feb 2004.
- J. Marchini and B. Howie. Genotype imputation for genome-wide association studies. Nature Reviews Genetics, 11(7):499–511, 2010.
- J. Marchini, B. Howie, S. Myers, G. McVean, and P. Donnelly. A new multipoint method for genome-wide association studies by imputation of genotypes. Nat. Genet., 39(7):906–913, Jul 2007.
- E. R. Mardis. The impact of next-generation sequencing technology on genetics. Trends Genet., 24(3):133–141, Mar 2008.

- P. Margaritte, C. Bonaiti-Pellie, M. C. King, and F. Clerget-Darpoux. Linkage of familial breast cancer to chromosome 17q21 may not be restricted to early-onset disease. Am. J. Hum. Genet., 50(6):1231–1234, Jun 1992.
- E. J. Masicampo and D. R. Lalande. A peculiar prevalence of p values just below .05. Q J Exp Psychol (Hove), Aug 2012.
- H. Matsuzaki, S. Dong, H. Loi, et al. Genotyping over 100,000 SNPs on a pair of oligonucleotide arrays. Nat. Methods, 1(2):109–111, Nov 2004a.
- H. Matsuzaki, H. Loi, S. Dong, et al. Parallel genotyping of over 10,000 SNPs using a one-primer assay on a high-density oligonucleotide array. Genome Res., 14(3):414–425, Mar 2004b.
- M. N. McCall, K. Uppal, H. A. Jaffee, M. J. Zilliox, and R. A. Irizarry. The Gene Expression Barcode: leveraging public data repositories to begin cataloging the human and murine transcriptomes. Nucleic Acids Res., 39(Database issue):D1011–1015, Jan 2011.
- M. I. McCarthy and E. Zeggini. Genome-wide association studies in type 2 diabetes. Curr. Diab. Rep., 9(2):164–171, Apr 2009.
- J. L. McCauley, R. L. Zuvich, A. L. Beecham, et al. Comprehensive follow-up of the first genome-wide association study of multiple sclerosis identifies KIF21B and TMEM39A as susceptibility loci. Hum. Mol. Genet., 19(5):953–962, Mar 2010.
- D. McGonagle, A. Aziz, L. J. Dickie, and M. F. McDermott. An integrated classification of pediatric inflammatory diseases, based on the concepts of autoinflammation and the immunological disease continuum. Pediatr. Res., 65(5 Pt 2):38R–45R, May 2009.
- A. McKenna, M. Hanna, E. Banks, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res., 20(9):1297–1303, Sep 2010.
- W. McLaren, B. Pritchard, D. Rios, et al. Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. Bioinformatics, 26(16):2069–2070, Aug 2010.
- G. A. McVean and N. J. Cardin. Approximating the coalescent with recombination. Philos. Trans. R. Soc. Lond., B, Biol. Sci., 360(1459):1387–1393, Jul 2005.
- G. A. McVean, S. R. Myers, S. Hunt, et al. The fine-scale structure of recombination rate variation in the human genome. Science, 304(5670):581–584, Apr 2004.

- R. Mead. A generalised logit-normal distribution. *Biometrics*, 21(3):721–732, 1965.
- G. Mendel. Versuche über pflanzenhybriden. *Verh. Naturforsch.*, 4, 1866.
- I. L. Mero, A. R. Lorentzen, M. Ban, et al. A rare variant of the TYK2 gene is confirmed to be associated with multiple sclerosis. *Eur. J. Hum. Genet.*, 18(4): 502–504, Apr 2010.
- Y. Miki, J. Swensen, D. Shattuck-Eidens, et al. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science*, 266(5182):66–71, Oct 1994.
- R. E. Mills, W. S. Pittard, J. M. Mullaney, et al. Natural genetic variation caused by small insertions and deletions in the human genome. *Genome Res.*, 21(6): 830–839, Jun 2011a.
- R. E. Mills, K. Walter, C. Stewart, et al. Mapping copy number variation by population-scale genome sequencing. *Nature*, 470(7332):59–65, Feb 2011b.
- Y. Minegishi, M. Saito, S. Tsuchiya, et al. Dominant-negative mutations in the DNA-binding domain of STAT3 cause hyper-IgE syndrome. *Nature*, 448(7157): 1058–1062, Aug 2007.
- A. Molven and P. R. Njølstad. Role of molecular genetics in transforming diagnosis of diabetes mellitus. *Expert Rev. Mol. Diagn.*, 11(3):313–320, Apr 2011.
- S.P. Morgan and J.D. Teachman. Logistic regression: Description, examples, and comparisons. *Journal of Marriage and Family*, 50(4):929–936, 1988.
- T. M. Morgan, H. M. Krumholz, R. P. Lifton, and J. A. Spertus. Nonvalidation of reported genetic risk factors for acute coronary syndrome in a large-scale replication study. *JAMA*, 297(14):1551–1561, Apr 2007.
- J. A. Morris, J. C. Randall, J. B. Maller, and J. C. Barrett. Evoker: a visualization tool for genotype intensity data. *Bioinformatics*, 26(14):1786–1787, Jul 2010.
- N. E. Morton. Sequential tests for the detection of linkage. *Am. J. Hum. Genet.*, 7(3):277–318, Sep 1955.
- J. Mudter, B. Weigmann, B. Bartsch, et al. Activation pattern of signal transducers and activators of transcription (STAT) factors in inflammatory bowel diseases. *Am. J. Gastroenterol.*, 100(1):64–72, Jan 2005.
- K. Mullis, F. Faloona, S. Scharf, et al. Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Cold Spring Harb. Symp. Quant. Biol.*, 51 Pt 1:263–273, 1986.

- R. M. Myers, J. Stamatoyannopoulos, M. Snyder, et al. A user's guide to the encyclopedia of DNA elements (ENCODE). *PLoS Biol.*, 9(4):e1001046, Apr 2011.
- R. P. Nair, K. C. Duffin, C. Helms, et al. Genome-wide scan reveals association of psoriasis with IL-23 and NF-kappaB pathways. *Nat. Genet.*, 41(2):199–204, Feb 2009.
- M. A. Nalls, V. Plagnol, D. G. Hernandez, et al. Imputation of sequence variants for identification of genetic risks for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet*, 377:641–649, 2011.
- B. M. Neale, Y. Kou, L. Liu, et al. Patterns and rates of exonic de novo mutations in autism spectrum disorders. *Nature*, 485(7397):242–245, May 2012.
- B. M. Neale, M. A. Rivas, B. F. Voight, et al. Testing for an unusual distribution of rare variants. *PLoS Genet.*, 7(3):e1001322, Mar 2011.
- M.C. Neale and L.R. Cardon. *Methodology for genetic studies of twins and families*. Number 67 in Series D: Behavioural and Social Sciences. Springer, 1992.
- A. C. Need, D. K. Attix, J. M. McEvoy, et al. Failure to replicate effect of Kibra on human memory in two large cohorts of European origin. *Am. J. Med. Genet. B Neuropsychiatr. Genet.*, 147B(5):667–668, Jul 2008.
- S. Nejentsev, J. M. Howson, N. M. Walker, et al. Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. *Nature*, 450(7171):887–892, Dec 2007.
- S. Nejentsev, N. Walker, D. Riches, M. Egholm, and J. A. Todd. Rare variants of IFIH1, a gene implicated in antiviral responses, protect against type 1 diabetes. *Science*, 324(5925):387–389, Apr 2009.
- J. A. Nelder and R. Mead. A simplex algorithm for function minimization. *Computer Journal*, 7:308–313, 1965.
- P. C. Ng, S. S. Murray, S. Levy, and J. C. Venter. An agenda for personalized medicine. *Nature*, 461(7265):724–726, Oct 2009.
- S. B. Ng, K. J. Buckingham, C. Lee, et al. Exome sequencing identifies the cause of a mendelian disorder. *Nat. Genet.*, 42(1):30–35, Jan 2010.
- NHLBI GO Exome Sequencing Project (ESP). *Exome Variant Server*, 2012. URL <http://evs.gs.washington.edu/EVS/>.

- R. Nielsen, J. S. Paul, A. Albrechtsen, and Y. S. Song. Genotype and SNP calling from next-generation sequencing data. Nat. Rev. Genet., 12(6):443–451, Jun 2011.
- Y. Nievergelt. How (not) to solve quadratic equations. The College Mathematics Journal, 34(2):90–104, 2003.
- L. Nistico, R. Buzzetti, L. E. Pritchard, et al. The CTLA-4 gene region of chromosome 2q33 is linked to, and associated with, type 1 diabetes. Belgian Diabetes Registry. Hum. Mol. Genet., 5(7):1075–1080, Jul 1996.
- L. D. Notarangelo, A. Fischer, R. S. Geha, et al. Primary immunodeficiencies: 2009 update. J. Allergy Clin. Immunol., 124(6):1161–1178, Dec 2009.
- M. C. O’Donovan, N. Craddock, N. Norton, et al. Identification of loci associated with schizophrenia by genome-wide association and follow-up. Nat. Genet., 40(9):1053–1055, Sep 2008.
- A. Papassotiropoulos, D. A. Stephan, M. J. Huentelman, et al. Common Kibra alleles are associated with human memory performance. Science, 314(5798):475–478, Oct 2006.
- M. Parkes, J. C. Barrett, N. J. Prescott, et al. Sequence variants in the autophagy gene IRGM and multiple other replicating loci contribute to Crohn’s disease susceptibility. Nat. Genet., 39(7):830–832, Jul 2007.
- S. Y. Patel, R. Doffinger, G. Barcenas-Morales, and D. S. Kumararatne. Genetically determined susceptibility to mycobacterial infection. J. Clin. Pathol., 61(9):1006–1012, Sep 2008.
- N. Patil, A. J. Berno, D. A. Hinds, et al. Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. Science, 294(5547):1719–1723, Nov 2001.
- L. Pauling, H. A. Itano, S.J. Singer, and I.C. Wells. Sickle cell anemia: a molecular disease. Science, 110(2865):543–548, Nov 1949.
- P. D. Pharoah, A. Antoniou, M. Bobrow, et al. Polygenic susceptibility to breast cancer and implications for prevention. Nat. Genet., 31(1):33–36, May 2002.
- J.K. Pickrell and J.K. Pritchard. Inference of population splits and mixtures from genome-wide allele frequency data. available from nature precedings <http://hdl.handle.net/10101/npre.2012.6956.1>. Nature proceedings, 2012.
- M. Pop. Genome assembly reborn: recent computational challenges. Brief. Bioinformatics, 10(4):354–366, Jul 2009.

- C. Power and J. Elliott. Cohort profile: 1958 British birth cohort (National Child Development Study). *International Journal of Epidemiology*, 35(1):34–41, 2006.
- A. L. Price, M. E. Weale, N. Patterson, et al. Long-range LD can confound genome scans in admixed populations. *Am. J. Hum. Genet.*, 83(1):132–135, Jul 2008.
- A.L. Price, N.J. Patterson, R.M. Plenge, et al. Principal components analysis corrects for stratification in genome-wide association studies. *Nature genetics*, 38(8):904–909, 2006.
- J. K. Pritchard. Are rare variants responsible for susceptibility to complex diseases? *Am. J. Hum. Genet.*, 69(1):124–137, Jul 2001.
- The 1000 Genomes Project. A map of human genome variation from population-scale sequencing. *Nature*, 467(7319):1061–1073, Oct 2010.
- The 1000 Genomes Project. An integrated map of genetic variation from 1,092 human genomes. *Under revision at Nature*, 2012.
- R.C. Punnett. Mendelism in relation to disease. *Proceedings of the Royal Society of Medicine*, 1:135, 1908.
- S. Purcell, B. Neale, K. Todd-Brown, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.*, 81(3):559–575, Sep 2007.
- S.M. Purcell, N.R. Wray, J.L. Stone, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*, 460(7256):748–752, 2009.
- H. Q. Qu, S. F. Grant, J. P. Bradfield, et al. Association of RASGRP1 with type 1 diabetes is revealed by combined follow-up of two genome-wide studies. *J. Med. Genet.*, 46(8):553–554, Aug 2009.
- M. Quail, M. E. Smith, P. Coupland, et al. A tale of three next generation sequencing platforms: comparison of Ion torrent, pacific biosciences and illumina MiSeq sequencers. *BMC Genomics*, 13(1):341, Jul 2012.
- M. Raabe. *Hemophilia*. Chelsea House Pub, 2008.
- S. Raychaudhuri, R. M. Plenge, E. J. Rossin, et al. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.*, 5(6):e1000534, Jun 2009a.
- S. Raychaudhuri, B. P. Thomson, E. F. Remmers, et al. Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk. *Nat. Genet.*, 41(12):1313–1318, Dec 2009b.

- L.J. Reed and J. Berkson. The application of the logistic function to experimental data. The Journal of Physical Chemistry, 33(5):760–779, 1929.
- T. Reich, JW James, and CA Morris. The use of multiple thresholds in determining the mode of transmission of semi-continuous traits*. Annals of human genetics, 36(2):163–184, 1972.
- J. D. Reveille, A. M. Sims, P. Danoy, et al. Genome-wide association study of ankylosing spondylitis identifies non-MHC susceptibility loci. Nat. Genet., 42(2):123–127, Feb 2010.
- F.V. Rijdsdijk and P.C. Sham. Analytic approaches to twin data using structural equation models. Briefings in bioinformatics, 3(2):119–133, 2002.
- J. R. Riordan, J. M. Rommens, B. Kerem, et al. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. Science, 245(4922):1066–1073, Sep 1989.
- J. Rios, E. Stein, J. Shendure, H.H. Hobbs, and J.C. Cohen. Identification by whole-genome resequencing of gene defect responsible for severe hypercholesterolemia. Hum. Mol. Genet., 19(22):4313–4318, 2010.
- N. Risch. Linkage strategies for genetically complex traits. I. Multilocus models. Am. J. Hum. Genet., 46(2):222–228, Feb 1990.
- N. Risch and K. Merikangas. The future of genetic studies of complex human diseases. Science, 273(5281):1516–1517, Sep 1996.
- M. A. Rivas, M. Beaudoin, A. Gardet, et al. Deep resequencing of GWAS loci identifies independent rare variants associated with inflammatory bowel disease. Nat. Genet., 43(11):1066–1073, Nov 2011.
- J. M. Rommens, M. C. Iannuzzi, B. Kerem, et al. Identification of the cystic fibrosis gene: chromosome walking and jumping. Science, 245(4922):1059–1065, Sep 1989.
- K. R. Rosenbloom, T. R. Dreszer, J. C. Long, et al. ENCODE whole-genome data in the UCSC Genome Browser: update 2012. Nucleic Acids Res., 40(Database issue):D912–917, Jan 2012.
- E. J. Rossin, K. Lage, S. Raychaudhuri, et al. Proteins encoded in genomic regions associated with immune-mediated disease physically interact and suggest underlying biology. PLoS Genet., 7(1):e1001273, 2011.
- D. Roskopf, A. Bornhorst, C. Rimbach, et al. Comment on “A common genetic variant is associated with adult and childhood obesity”. Science, 315(5809):187; author reply 187, Jan 2007.

- M. Ruffalo, T. LaFramboise, and M. Koyuturk. Comparative analysis of algorithms for next-generation sequencing read alignment. Bioinformatics, 27(20):2790–2796, Oct 2011.
- R.K. Russell and J. Satsangi. IBD: a family affair. Best Practice & Research Clinical Gastroenterology, 18(3):525–539, 2004.
- R. Sachidanandam, D. Weissman, S. C. Schmidt, et al. A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. Nature, 409(6822):928–933, Feb 2001.
- D.L. Sackett, J.J. Deeks, and D.G. Altman. Down with odds ratios! Evidence Based Medicine, 1(6):164–166, 1996.
- N. J. Samani, J. Erdmann, A. S. Hall, et al. Genomewide association analysis of coronary artery disease. N. Engl. J. Med., 357(5):443–453, Aug 2007.
- F. Sanger, S. Nicklen, and A. R. Coulson. DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. U.S.A., 74(12):5463–5467, Dec 1977.
- J. Satsangi, D. P. Jewell, and J. I. Bell. The genetics of inflammatory bowel disease. Gut, 40(5):572–574, May 1997.
- S. Sawcer. The genetic aspects of multiple sclerosis. Ann Indian Acad Neurol, 12(4):206–214, Oct 2009.
- S. Sawcer, M. Ban, J. Wason, and F. Dudbridge. What role for genetics in the prediction of multiple sclerosis? Ann. Neurol., 67(1):3–10, Jan 2010.
- S. Sawcer, G. Hellenthal, M. Pirinen, et al. Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. Nature, 476(7359):214–219, Aug 2011.
- S. Sawcer and J. Wason. Risk in complex genetics: "All models are wrong but some are useful". Ann Neurol, Apr 2012.
- E. E. Schadt, C. Molony, E. Chudin, et al. Mapping the genetic architecture of gene expression in human liver. PLoS Biol., 6(5):e107, May 2008.
- K. Schaper, H. Kolsch, J. Popp, M. Wagner, and F. Jessen. KIBRA gene variants are associated with episodic memory in healthy elderly. Neurobiol. Aging, 29(7):1123–1125, Jul 2008.
- P. Scheet and M. Stephens. A fast and flexible statistical model for large-scale population genotype data: applications to inferring missing genotypes and haplotypic phase. Am. J. Hum. Genet., 78(4):629–644, Apr 2006.

- N.R. Schneider, WR Williams, RSK Chaganti, et al. Genetic epidemiology of familial aggregation of cancer. Advances in cancer research, 47:1–36, 1986.
- H. Schunkert, I. R. Konig, S. Kathiresan, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. Nat. Genet., 43: 333–338, 2011.
- L.J. Scott, P. Muglia, X.Q. Kong, et al. Genome-wide association and meta-analysis of bipolar disorder in individuals of European ancestry. Proc. Natl. Acad. Sci. U.S.A., 106(18):7501–7506, 2009.
- R. A. Scott, V. Lagou, R. P. Welch, et al. Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. Nat. Genet., Aug 2012.
- J.M. Seddon, R. Reynolds, J. Maller, et al. Prediction model for prevalence and incidence of advanced age-related macular degeneration based on genetic, demographic, and environmental variables. Invest. Ophth. Vis. Sci., 50(5):2044–2053, 2009.
- B. Servin and M. Stephens. Imputation-based analysis of association studies: candidate regions and quantitative traits. PLoS Genet., 3(7):e114, Jul 2007.
- E. Sezgin, J. M. Lind, S. Shrestha, et al. Association of Y chromosome haplogroup I with HIV progression, and HAART outcome. Hum. Genet., 125(3):281–294, Apr 2009.
- T. S. Shah, J. Z. Liu, J. A. Floyd, et al. optiCall: a robust genotype-calling algorithm for rare, low-frequency and common variants. Bioinformatics, 28(12): 1598–1603, Jun 2012.
- M. H. Shaw, N. Kamada, N. Warner, Y. G. Kim, and G. Nunez. The ever-expanding function of NOD2: autophagy, viral recognition, and T cell activation. Trends Immunol., 32(2):73–79, Feb 2011.
- P. Sherlock, B. M. Bell, H. Steinberg, and T. P. Almy. Familial occurrence of regional enteritis and ulcerative colitis. Gastroenterology, 45:413–420, Sep 1963.
- S. T. Sherry, M. Ward, and K. Sirotkin. dbSNP-database for single nucleotide polymorphisms and other classes of minor genetic variation. Genome Res., 9 (8):677–679, Aug 1999.
- S. Shivananda, J. Lennard-Jones, R. Logan, et al. Incidence of inflammatory bowel disease across Europe: is there a difference between north and south? Results of the European Collaborative Study on Inflammatory Bowel Disease (EC-IBD). Gut, 39(5):690–697, Nov 1996.

- S.I. Shyn, J. Shi, J.B. Kraft, et al. Novel loci for major depression identified by genome-wide association study of Sequenced Treatment Alternatives to Relieve Depression and meta-analysis of three studies. Mol. Psychiatr., 16(2):202–215, 2009.
- A. Siepel, G. Bejerano, J. S. Pedersen, et al. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res., 15(8):1034–1050, Aug 2005.
- M. S. Silverberg, J. H. Cho, J. D. Rioux, et al. Ulcerative colitis-risk loci on chromosomes 1p36 and 12q15 found by genome-wide association study. Nat. Genet., 41(2):216–220, Feb 2009.
- L. M. Smith, J. Z. Sanders, R. J. Kaiser, et al. Fluorescence detection in automated DNA sequence analysis. Nature, 321(6071):674–679, 1986.
- D. J. Smyth, V. Plagnol, N. M. Walker, et al. Shared and distinct genetic variants in type 1 diabetes and celiac disease. N. Engl. J. Med., 359(26):2767–2777, Dec 2008.
- H. Song, S. J. Ramus, J. Tyrer, et al. A genome-wide association study identifies a new ovarian cancer susceptibility locus on 9p22.2. Nat. Genet., 41:996–1000, 2009.
- E. M. Southern. Detection of specific sequences among DNA fragments separated by gel electrophoresis. J. Mol. Biol., 98(3):503–517, Nov 1975.
- P. H. St George-Hyslop, R. E. Tanzi, R. J. Polinsky, et al. The genetic defect causing familial Alzheimer’s disease maps on chromosome 21. Science, 235(4791):885–890, Feb 1987.
- E. A. Stahl, S. Raychaudhuri, E. F. Remmers, et al. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. Nat. Genet., 42(6):508–514, Jun 2010.
- M. Stephens, N. J. Smith, and P. Donnelly. A new statistical method for haplotype reconstruction from population data. Am. J. Hum. Genet., 68(4):978–989, Apr 2001.
- A. Sterner-Kock, I. S. Thorey, K. Koli, et al. Disruption of the gene encoding the latent transforming growth factor-beta binding protein 4 (LTBP-4) causes abnormal lung development, cardiomyopathy, and colorectal cancer. Genes Dev., 16(17):2264–2273, Sep 2002.
- R. Stiratelli, N. Laird, and J. H. Ware. Random-effects models for serial observations with binary response. Biometrics, 40(4):961–971, Dec 1984.

- A. Strange, F. Capon, C. C. Spencer, et al. A genome-wide association study identifies new psoriasis susceptibility loci and an interaction between HLA-C and ERAP1. *Nat. Genet.*, 42(11):985–990, Nov 2010.
- P. E. Stuart, R. P. Nair, E. Ellinghaus, et al. Genome-wide association analysis identifies three psoriasis susceptibility loci. *Nat. Genet.*, 42(11):1000–1004, Nov 2010.
- Y. J. Sung, L. Wang, T. Rankinen, C. Bouchard, and D. C. Rao. Performance of genotype imputations using data from the 1000 Genomes Project. *Hum. Hered.*, 73(1):18–25, 2012.
- I. Surolia, S. P. Pirnie, V. Chellappa, et al. Functionally defective germline variants of sialic acid acetyltransferase in autoimmunity. *Nature*, 466(7303):243–247, Jul 2010.
- A. D. Swafford, J. M. Howson, L. J. Davison, et al. An allele of IKZF1 (Ikaros) conferring susceptibility to childhood acute lymphoblastic leukemia protects against type 1 diabetes. *Diabetes*, 60(3):1041–1044, Mar 2011.
- A. C. Syvanen. Toward genome-wide SNP genotyping. *Nat. Genet.*, 37 Suppl: 5–10, Jun 2005.
- Y. Y. Teo, A. E. Fry, K. Bhattacharya, et al. Genome-wide comparisons of variation in linkage disequilibrium. *Genome Res.*, 19(10):1849–1860, Oct 2009.
- Y. Y. Teo, M. Inouye, K. S. Small, et al. A genotype calling algorithm for the Illumina BeadArray platform. *Bioinformatics*, 23(20):2741–2746, Oct 2007.
- Y. Y. Teo, K. S. Small, and D. P. Kwiatkowski. Methodological challenges of genome-wide association analysis in Africa. *Nat. Rev. Genet.*, 11(2):149–160, Feb 2010.
- The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature*, 489:57–74, Sep 2012.
- The Huntington’s Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington’s disease chromosomes. . *Cell*, 72(6):971–983, Mar 1993.
- The MalariaGEN Consortium. Genome-wide and fine-resolution association analysis of malaria in West Africa. *Nat. Genet.*, 41(6):657–665, Jun 2009.
- W. Thomson, A. Barton, X. Ke, et al. Rheumatoid arthritis association at 6q23. *Nature genetics*, 39(12):1431–1433, 2007.

- T. Thye, E. Owusu-Dabo, F. O. Vannberg, et al. Common variants at 11p13 are associated with susceptibility to tuberculosis. *Nat. Genet.*, 44(3):257–259, Mar 2012.
- J. A. Todd, J. I. Bell, and H. O. McDevitt. HLA-DQ beta gene contributes to susceptibility and resistance to insulin-dependent diabetes mellitus. *Nature*, 329(6140):599–604, 1987.
- J. A. Todd, N. M. Walker, J. D. Cooper, et al. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat. Genet.*, 39(7):857–864, Jul 2007.
- S. P. Tsai, R. J. Hardy, and C. P. Wen. The standardized mortality ratio and life expectancy. *Am. J. Epidemiol.*, 135(7):824–831, Apr 1992.
- L. C. Tsui, M. Buchwald, D. Barker, et al. Cystic fibrosis locus defined by a genetically linked polymorphic DNA marker. *Science*, 230(4729):1054–1057, Nov 1985.
- C. Turnbull, S. Ahmed, J. Morrison, et al. Genome-wide association study identifies five new breast cancer susceptibility loci. *Nat. Genet.*, 42(6):504–517, 2010.
- D. A. van Heel, S. A. Fisher, A. Kirby, et al. Inflammatory bowel disease susceptibility loci defined by genome scan meta-analysis of 1952 affected relative pairs. *Hum. Mol. Genet.*, 13(7):763–770, Apr 2004.
- B. Vanhaesebroeck, J. Guillermet-Guibert, M. Graupera, and B. Bilanges. The emerging mechanisms of isoform-specific PI3K signalling. *Nat. Rev. Mol. Cell Biol.*, 11(5):329–341, May 2010.
- M.H. Vatn. Environmental factors in the epidemiology of inflammatory bowel disease. *Crohn's Disease and Ulcerative Colitis: From Epidemiology and Immunobiology to a Rational Diagnostic and Therapeutic Approach*, page 17, 2011.
- J. C. Venter, M. D. Adams, E. W. Myers, et al. The sequence of the human genome. *Science*, 291(5507):1304–1351, Feb 2001.
- P.F. Verhulst. Notice sur la loi que la population suit dans son accroissement. *Correspondance Mathématique et Physique*, publiée par A. Quetelet, 10:113–121, 1838.
- B. F. Voight, G. M. Peloso, M. Orho-Melander, et al. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet*, 380(9841):572–580, Aug 2012.

- B. F. Voight, L. J. Scott, V. Steinthorsdottir, et al. Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nat. Genet.*, 42:579–589, 2010.
- S. Wacholder, P. Hartge, R. Prentice, et al. Performance of common genetic variants in breast-cancer risk models. *New Engl. J. Med.*, 362(11):986–993, 2010.
- B. J. Wainwright, P. J. Scambler, J. Schmidtke, et al. Localization of cystic fibrosis locus to human chromosome 7cen-q22. *Nature*, 318(6044):384–385, 1985.
- C. Wallace, D. J. Smyth, M. Maisuria-Armer, et al. The imprinted DLK1-MEG3 gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. *Nat. Genet.*, 42(1):68–71, Jan 2010.
- D. G. Wang, J. B. Fan, C. J. Siao, et al. Large-scale identification, mapping, and genotyping of single-nucleotide polymorphisms in the human genome. *Science*, 280(5366):1077–1082, May 1998.
- K. Wang, M. Li, and H. Hakonarson. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.*, 38(16):e164, Sep 2010.
- Y. H. Wang and Y. J. Liu. OX40-OX40L interactions: a promising therapeutic target for allergic diseases? *J. Clin. Invest.*, 117(12):3655–3657, Dec 2007.
- J. D. Watson and F. H. Crick. Molecular structure of nucleic acids; a structure for deoxyribose nucleic acid. *Nature*, 171(4356):737–738, Apr 1953.
- J. L. Weber and P. E. May. Abundant class of human DNA polymorphisms which can be typed using the polymerase chain reaction. *Am. J. Hum. Genet.*, 44(3):388–396, Mar 1989.
- Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*, 447(7145):661–678, Jun 2007.
- K. A. Wetterstrand. DNA sequencing costs: data from the NHGRI large-scale genome sequencing program. Retrieved 16/08/2012, Available at: www.genome.gov/sequencingcosts. Accessed [2011-10-25], 2012.
- C. C. Whitacre. Sex differences in autoimmune disease. *Nat. Immunol.*, 2(9):777–780, Sep 2001.
- R. White, S. Woodward, M. Leppert, et al. A closely linked genetic marker for cystic fibrosis. *Nature*, 318(6044):382–384, 1985.

- C. N. Williams, K. Kocher, E. S. Lander, M. J. Daly, and J. D. Rioux. Using a genome-wide scan and meta-analysis to identify a novel IBD locus and confirm previously identified IBD loci. *Inflamm. Bowel Dis.*, 8(6):375–381, Nov 2002.
- L. H. Wise, J. S. Lanchbury, and C. M. Lewis. Meta-analysis of genome searches. *Ann. Hum. Genet.*, 63(Pt 3):263–272, May 1999.
- M. G. Wolfs, M. H. Hofker, C. Wijmenga, and T. W. van Haeften. Type 2 Diabetes Mellitus: New Genetic Insights will Lead to New Therapeutics. *Curr. Genomics*, 10(2):110–118, Apr 2009.
- K. Wong, T. M. Keane, J. Stalker, and D. J. Adams. Enhanced structural variant and breakpoint detection using SVMerge by integration of multiple detection methods and local assembly. *Genome Biol.*, 11(12):R128, 2010.
- E. A. Worthey, A. N. Mayer, G. D. Syverson, et al. Making a definitive diagnosis: successful clinical application of whole exome sequencing in a child with intractable inflammatory bowel disease. *Genet. Med.*, 13(3):255–262, Mar 2011a.
- E.A. Worthey, A.N. Mayer, G.D. Syverson, et al. Making a definitive diagnosis: successful clinical application of whole exome sequencing in a child with intractable inflammatory bowel disease. *Genet. Med.*, 13(3):255–262, 2011b.
- N. R. Wray and M. E. Goddard. Multi-locus models of genetic risk of disease. *Genome Med*, 2(2):10, 2010.
- N. R. Wray, M. E. Goddard, and P. M. Visscher. Prediction of individual genetic risk to disease from genome-wide association studies. *Genome Res.*, 17(10):1520–1528, Oct 2007.
- N. R. Wray, J. Yang, M. E. Goddard, and P. M. Visscher. The genetic interpretation of area under the ROC curve in genomic profiling. *PLoS Genet.*, 6(2):e1000864, Feb 2010.
- C. F. Wright and S. Gregory-Jones. Size of the direct-to-consumer genomic testing market. *Genet. Med.*, 12(9):594, Sep 2010.
- J. Yang, P. M. Visscher, and N. R. Wray. Sporadic cases are the norm for complex disease. *Eur. J. Hum. Genet.*, 18(9):1039–1043, Sep 2010.
- S. Yazdanyar, M. Weischer, and B. G. Nordestgaard. Genotyping for NOD2 genetic variants and crohn disease: a metaanalysis. *Clin. Chem.*, 55:1950–1957, 2009.
- S. Yoon, Z. Xuan, V. Makarov, K. Ye, and J. Sebat. Sensitive and accurate detection of copy number variants using read depth of coverage. *Genome Res.*, 19(9):1586–1592, Sep 2009.

- A. P. Yu, L. A. Cabanilla, E. Q. Wu, P. M. Mulani, and J. Chao. The costs of Crohn's disease in the United States and other Western countries: a systematic review. *Curr Med Res Opin*, 24(2):319–328, Feb 2008.
- E. Zeggini, L. J. Scott, R. Saxena, et al. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat. Genet.*, 40(5):638–645, May 2008.
- E. Zeggini, M. N. Weedon, C. M. Lindgren, et al. Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science*, 316(5829):1336–1341, Jun 2007.
- T. Zeller, P. Wild, S. Szymczak, et al. Genetics and beyond—the transcriptome of human monocytes and disease susceptibility. *PLoS ONE*, 5(5):e10693, 2010.
- D. R. Zerbino and E. Birney. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.*, 18(5):821–829, May 2008.
- B. Zhang and S. Horvath. A general framework for weighted gene co-expression network analysis. *Stat Appl Genet Mol Biol*, 4:Article17, 2005.
- F. Zhang, H. Liu, S. Chen, et al. Identification of two new loci at IL23R and RAB32 that influence susceptibility to leprosy. *Nature genetics*, 43(12):1247–51, 2011.
- H. Zhang, D. Massey, M. Tremelling, and M. Parkes. Genetics of inflammatory bowel disease: clues to pathogenesis. *Br. Med. Bull.*, 87:17–30, 2008.
- J. Zhang, L. Feuk, G. E. Duggan, R. Khaja, and S. W. Scherer. Development of bioinformatics resources for display and analysis of copy number and other structural variants in the human genome. *Cytogenet. Genome Res.*, 115(3-4): 205–214, 2006.
- X. J. Zhang, W. Huang, S. Yang, et al. Psoriasis genome-wide association study identifies susceptibility variants within LCE gene cluster at 1q21. *Nat. Genet.*, 41(2):205–210, Feb 2009.
- J. Zhu, M. C. Wiener, C. Zhang, et al. Increasing the power to detect causal associations by combining genotypic and expression data in segregating populations. *PLoS Comput. Biol.*, 3(4):e69, Apr 2007.
- O. Zuk, E. Hechter, S. R. Sunyaev, and E. S. Lander. The mystery of missing heritability: Genetic interactions create phantom heritability. *Proc. Natl. Acad. Sci. U.S.A.*, 109(4):1193–1198, Jan 2012.